

Polynomial Approximations ⁷

4.3.96

$$0 \leq x \leq \frac{\pi}{2}$$

$$\frac{\sin x}{x} = 1 + a_2 x^2 + a_4 x^4 + \epsilon(x)$$

$$|\epsilon(x)| \leq 2 \times 10^{-4}$$

$$a_2 = -.16605 \quad a_4 = .00761$$

4.3.97

$$0 \leq x \leq \frac{\pi}{2}$$

$$\frac{\sin x}{x} = 1 + a_2 x^2 + a_4 x^4 + a_6 x^6 + a_8 x^8 + a_{10} x^{10} + \epsilon(x)$$

$$|\epsilon(x)| \leq 2 \times 10^{-9}$$

$$a_2 = -.16666 \ 66664 \quad a_8 = .00000 \ 27526$$

$$a_4 = .00833 \ 33315 \quad a_{10} = -.00000 \ 00239$$

$$a_6 = -.00019 \ 84090$$

4.3.98

$$0 \leq x \leq \frac{\pi}{2}$$

$$\cos x = 1 + a_2 x^2 + a_4 x^4 + \epsilon(x)$$

$$|\epsilon(x)| \leq 9 \times 10^{-4}$$

$$a_2 = -.49670 \quad a_4 = .03705$$

4.3.99

$$0 \leq x \leq \frac{\pi}{2}$$

$$\cos x = 1 + a_2 x^2 + a_4 x^4 + a_6 x^6 + a_8 x^8 + a_{10} x^{10} + \epsilon(x)$$

$$|\epsilon(x)| \leq 2 \times 10^{-9}$$

$$a_2 = -.49999 \ 99963 \quad a_8 = .00002 \ 47609$$

$$a_4 = .04166 \ 66418 \quad a_{10} = -.00000 \ 02605$$

$$a_6 = -.00138 \ 88397$$

4.3.100

$$0 \leq x \leq \frac{\pi}{4}$$

$$\frac{\tan x}{x} = 1 + a_2 x^2 + a_4 x^4 + \epsilon(x)$$

$$|\epsilon(x)| \leq 1 \times 10^{-3}$$

$$a_2 = .31755 \quad a_4 = .20330$$

4.3.101

$$0 \leq x \leq \frac{\pi}{4}$$

$$\frac{\tan x}{x} = 1 + a_2 x^2 + a_4 x^4 + a_6 x^6 + a_8 x^8 + a_{10} x^{10} + a_{12} x^{12} + \epsilon(x)$$

$$|\epsilon(x)| \leq 2 \times 10^{-8}$$

$$a_2 = .33333 \ 14036 \quad a_8 = .02456 \ 50893$$

$$a_4 = .13339 \ 23995 \quad a_{10} = .00290 \ 05250$$

$$a_6 = .05337 \ 40603 \quad a_{12} = .00951 \ 68091$$

4.3.102

$$0 \leq x \leq \frac{\pi}{4}$$

$$* \quad x \cot x = 1 + a_2 x^2 + a_4 x^4 + \epsilon(x)$$

$$|\epsilon(x)| \leq 3 \times 10^{-5}$$

$$a_2 = -.332867 \quad a_4 = -.024369$$

4.3.103

$$0 \leq x \leq \frac{\pi}{4}$$

$$x \cot x = 1 + a_2 x^2 + a_4 x^4 + a_6 x^6 + a_8 x^8 + a_{10} x^{10} + \epsilon(x)$$

$$|\epsilon(x)| \leq 4 \times 10^{-10}$$

$$a_2 = -.33333 \ 33410 \quad a_8 = -.00020 \ 78504$$

$$a_4 = -.02222 \ 20287 \quad a_{10} = -.00002 \ 62619$$

$$a_6 = -.00211 \ 77168$$

Approximations in Terms of Chebyshev Polynomials ⁸

4.3.104

$$-1 \leq x \leq 1$$

$$T_n^*(x) = \cos n\theta, \cos \theta = 2x - 1 \quad (\text{see chapter 22})$$

$$\sin \frac{1}{2}\pi x = x \sum_{n=0}^{\infty} A_n T_n^*(x^2) \quad \cos \frac{1}{2}\pi x = \sum_{n=0}^{\infty} A_n T_n^*(x^2)$$

n	A_n	n	A_n
0	1.27627 8962	0	.47200 1216
1	-.28526 1569	1	-.49940 3258
2	.00911 8016	2	.02799 2080
3	-.00013 6587	3	-.00059 6695
4	.00000 1185	4	.00000 6704
5	-.00000 0007	5	-.00000 0047

⁷ The approximations 4.3.96 to 4.3.103 are from B. Carlson, M. Goldstein, Rational approximation of functions, Los Alamos Scientific Laboratory LA-1943, Los Alamos, N. Mex., 1955 (with permission).

⁸ The approximations 4.3.104 are from C. W. Clenshaw, Polynomial approximations to elementary functions, Math. Tables Aids Comp. 8, 143-147 (1954) (with permission).

*See page II.